Bosonization: Worksheet 5

Boson field operators

Christophe De Beule (christophe.debeule@gmail.com)

Starting from this exercise sheet, we use the following conventions. The density operator ρ_{qs} and corresponding commutator are defined as

$$\rho_{qs} = \sum_{k} c_{k+qs}^{\dagger} c_{ks}, \qquad [\rho_{qs}, \rho_{q's'}^{\dagger}] = -\delta_{ss'} \delta_{qq'} \epsilon_s \frac{qL}{2\pi},$$

Note that using c_{k-qs}^{\dagger} instead of c_{k+qs}^{\dagger} gives an extra minus sign in the commutator.

1 Boson field operators

Similar to the fermion field operators, we can define the boson field

$$\varphi(x) = \frac{\sqrt{\pi}}{L} (N_R + N_L) x + \frac{i\sqrt{\pi}}{L} \sum_{q \neq 0} \frac{1}{q} e^{-iqx - a|q|/2} (\rho_{qR} + \rho_{qL}),$$

where a > 0 is an infinitesimal regularization parameter. The boson field $\varphi(x)$ is chosen such that

$$\frac{1}{\sqrt{\pi}}\partial_x \varphi(x) = \rho_R(x) + \rho_L(x),$$

for $a \to 0$ and where $\rho_s(x) =: \psi_s^{\dagger}(x) \psi_s(x)$: with $\psi_s(x) = \frac{1}{\sqrt{L}} \sum_k e^{ikx} c_{ks}$ and $N_s = \sum_k : c_{ks}^{\dagger} c_{ks}$:

- (1) Show that the equation for $\partial_x \varphi(x)$ is correct and also show that $\varphi(x)$ is Hermitian.
- (2) Define another boson field, denoted as $\vartheta(x)^1$, in a similar manner as we defined $\varphi(x)$ and such that for $a \to 0$

$$\frac{1}{\sqrt{\pi}}\partial_x \vartheta(x) = \rho_R(x) - \rho_L(x).$$

(3) Show that

$$[\varphi(x), \vartheta(x')] = \frac{1}{2\pi} \left(\ln \left[1 - e^{i\frac{2\pi}{L}(x'-x+ia)} \right] - \ln \left[1 - e^{-i\frac{2\pi}{L}(x'-x-ia)} \right] \right).$$

Hint: $\sum_{n=1}^{\infty} z^n / n = -\ln(1-z)$ for |z| < 1.

(4) Show that

$$[\varphi(x), \vartheta(x')] \xrightarrow{L \to \infty, a \to 0} \begin{cases} \frac{i}{2} \operatorname{sgn}(x - x') & x \neq x', \\ 0 & x = x'. \end{cases}$$

Hint: Keep terms in 1/L and note that the complex logarithm is defined as $\ln z \equiv \ln |z| + i \arg z$ for $z \neq 0$ and $\arg z \in (-\pi, \pi]$, i.e. we put the branch cut along the negative real axis.

(5) Use this result to show that

$$[\varphi(x), \partial_{x'}\vartheta(x')] \xrightarrow{L\to\infty, a\to 0} -i\delta(x-x').$$

Note that if we define $\Pi(x) = -\partial_x \vartheta(x)$, we obtain the canonical form $[\varphi(x), \Pi(x')] = i\delta(x-x')$. Hint: $\delta(x) = \partial_x \Theta(x)$.

¹The boson field $\vartheta(x)$ is not to be confused with the step function $\Theta(x)$.

(6) Now start from (3) and take the derivative with respect to x'. Show that

$$[\varphi(x), \partial_{x'}\vartheta(x')] = -i\left[\frac{a/\pi}{(x'-x)^2 + a^2} - \frac{1}{L}\right] + \mathcal{O}(a/L^2),$$

and show that you obtain the same result as (5) in the limits $L \to \infty$ and $a \to 0$.

(7) Integrate your result from (6) with respect to x' to find

$$[\varphi(x), \vartheta(x')] = C + \frac{i}{\pi} \arctan\left(\frac{x - x'}{a}\right) + \frac{ix'}{L},$$

where we only kept the lowest-order terms and C is an integration constant. Determine C by requiring that $[\varphi(x), \vartheta(x)] = 0$. Finally take the limits $L \to \infty$ and $a \to 0$ to obtain the result in (4). Note that for finite a and L, the sign function is smeared over a range of order a and that there is a term of order 1/L.

2 Free Hamiltonian in terms of boson fields

Remember that the free Hamiltonian could be written as

$$H_0 = \frac{\pi v_F}{L} \sum_{s} \sum_{q \neq 0} : \rho_{-qs} \rho_{qs} : + \frac{\pi v_F}{L} \sum_{s} N_s^2,$$

where we put $\hbar = 1$. Note that the second term is different than in worksheet 3. This is because we are now using anti-periodic boundary conditions instead of periodic boundary conditions. The type of boundary condition makes no difference in the limit $L \to \infty$.

(1) Show that

$$\frac{1}{2} \int_{-L/2}^{L/2} dx \, (\partial_x \varphi(x))^2 = \frac{\pi}{2L} \sum_{q \neq 0} \left(\rho_{-qR} + \rho_{-qL} \right) \left(\rho_{qR} + \rho_{qL} \right) + \frac{\pi}{2L} \left(N_R + N_L \right)^2,$$

for $a \to 0$. Hint: Note that $q = \frac{2\pi}{L} n$ $(n \in \mathbb{Z})$, independent of the boundary conditions.

(2) Perform the same calculation for the boson field $\vartheta(x)$ and show that

$$H_0 = \frac{v_F}{2} \int_{-L/2}^{L/2} dx : \left[\left(\partial_x \varphi(x) \right)^2 + \left(\partial_x \vartheta(x) \right)^2 \right] :,$$

which only holds for anti-periodic boundary conditions if we do not take the limit $L \to \infty$.

(3) Finally, show that H_0 can also be written as

$$H_0 = \pi v_F \int_{-L/2}^{L/2} dx : [\rho_R^2(x) + \rho_L^2(x)] : .$$

We need normal ordering in the expressions above, because even though $\partial_x \varphi(x)$ and $\partial_x \vartheta(x)$ are normal ordered, their squares are not in general. Indeed, $\langle :A: \rangle_0 = 0$ but $\langle :A: A: \rangle_0 = \langle A^2 \rangle_0 - (\langle A \rangle_0)^2$.